

Symmetric Nonnegative 5×5 Matrices Realizing Previously Unknown Region

Oren Spector

March 25, 2014

Abstract

In this paper we present some symmetric nonnegative 5×5 matrix families that realize a previously unknown region. We also prove that these and other symmetric nonnegative 5×5 matrix families are closed under perturbations first presented in W. Guo. Eigenvalues of nonnegative matrices. *Linear Algebra and its Applications*, 266:261–270, 1997.

1 Introduction

Let $\sigma = (\lambda_1, \lambda_2, \dots, \lambda_n)$ be a list of complex numbers. The problem of determining necessary and sufficient conditions for σ to be the eigenvalues of a nonnegative $n \times n$ matrix is called the nonnegative inverse eigenvalue problem (NIEP). When σ is a list of real numbers the problem of determining necessary and sufficient conditions for σ to be the eigenvalues of a nonnegative $n \times n$ matrix is called the real nonnegative inverse eigenvalue problem (RNIEP). When σ is a list of real numbers the problem of determining necessary and sufficient conditions for σ to be the eigenvalues of a symmetric nonnegative $n \times n$ matrix is called the symmetric nonnegative inverse eigenvalue problem (SNIEP). All three problems are currently unsolved for $n \geq 5$.

Loewy and London [7] have solved NIEP for $n = 3$ and RNIEP for $n = 4$. Moreover, RNIEP and SNIEP are the same for $n \leq 4$. This can be seen from papers by Fiedler [3] and Loewy and London [7]. However, it has been shown by Johnson et al. [5] that RNIEP and SNIEP are different in general. More results about the general NIEP, RNIEP and SNIEP can be found in [1]. Other results about SNIEP for $n = 5$ can be found in [9, 8, 2, 11].

Guo [4] proved the realizability of some real eigenvalue perturbations of a realizable spectrum in NIEP and RNIEP. Laffey [6] extended this result to some non-real eigenvalue perturbations. Rojo and Soto [10] extended this result to symmetric nonnegative circulant matrices.

This paper is organized as follows: In Section 2 we present some notations and results concerning SNIEP. In Section 3 we present some matrix patterns and their properties. In Section 4 we show that these matrix patterns realize a previously unknown region. In Section 5 we prove Guo-type results for several families of 5×5 symmetric nonnegative matrices, including those presented in Section 3. Finally, in Section 6 we show some examples that illustrate our results.

2 Preliminaries and Notations

By the Perron-Frobenius Theorem we know that if σ is the spectrum of a non-negative matrix A then

$$\rho = \max_{\lambda \in \sigma} |\lambda| \in \sigma. \quad (1)$$

We call ρ the Perron eigenvalue of A . Also, by the nonnegativity of A we have

$$\sum_{\lambda \in \sigma} \lambda \geq 0. \quad (2)$$

The following result is due to Fiedler [3], who extended a result due to Suleimanova [12]:

Theorem 1. *Let $\lambda_1 \geq 0 \geq \lambda_2 \geq \dots \geq \lambda_n$ and $\sum_{i=1}^n \lambda_i \geq 0$. Then there exists a symmetric nonnegative $n \times n$ matrix with a spectrum $\sigma = (\lambda_1, \lambda_2, \dots, \lambda_n)$, where λ_1 is its Perron eigenvalue.*

By [7] and [3] we have the following theorem:

Theorem 2. *Let $\sigma = (\lambda_1, \lambda_2, \lambda_3, \lambda_4)$ be a list of real numbers. Necessary and sufficient conditions for σ to be the spectrum of a 4×4 symmetric nonnegative matrix are (1) and (2).*

The following necessary condition due to McDonald and Neumann [9] was extended by Loewy and McDonald [8]:

Lemma 1. *Let A be a 5×5 nonnegative symmetric matrix with a spectrum $\sigma = (\lambda_1, \lambda_2, \dots, \lambda_5)$ such that $\lambda_1 \geq \lambda_2 \geq \lambda_3 \geq \lambda_4 \geq \lambda_5$. Then*

$$\lambda_1 + \lambda_3 + \lambda_4 \geq 0. \quad (3)$$

Lemma 2. *Let a list of real numbers $\sigma = (\lambda_1, \lambda_2, \dots, \lambda_5)$ satisfy $\lambda_1 \geq \lambda_2 > 0 \geq \lambda_3 \geq \lambda_4 \geq \lambda_5$. Necessary and sufficient conditions for σ to be the spectrum of a 5×5 symmetric nonnegative matrix are (1), (2) and (3).*

Proof. Let A be a 5×5 symmetric nonnegative matrix with a spectrum $\sigma = (\lambda_1, \lambda_2, \dots, \lambda_5)$ such that $\lambda_1 \geq \lambda_2 \geq \lambda_3 \geq \lambda_4 \geq \lambda_5$. Therefore, (1) and (2) are met. By Lemma 1 condition (3) is met as well.

Let a list of real numbers $\sigma = (\lambda_1, \lambda_2, \dots, \lambda_5)$ satisfy $\lambda_1 \geq \lambda_2 > 0 \geq \lambda_3 \geq \lambda_4 \geq \lambda_5$, (1), (2) and (3). Realizing 5×5 symmetric nonnegative matrices for this case are given in [2]. $\square$

We denote the spectrum of a 5×5 nonnegative real symmetric matrix by the list of real numbers $\sigma = (\lambda_1, \lambda_2, \dots, \lambda_5)$. Unless otherwise stated we assume that

$$\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_5 \geq -\lambda_1. \quad (4)$$

Let $\alpha \in \mathbb{R}$. We define $\alpha \cdot \sigma = (\alpha\lambda_1, \alpha\lambda_2, \dots, \alpha\lambda_5)$.

By [8] we know that the only unknown realizable region of SNIEP for $n = 5$ is when $\lambda_3 > \sum_{i=1}^5 \lambda_i$.

3 Matrix Patterns

In this section we define several matrix patterns and calculate their spectra. Let $\sigma = (\lambda_1, \lambda_2, \dots, \lambda_5)$ be a list of complex numbers and let $g \in \mathbb{C}$. Define

$$A(\sigma) = \frac{1}{2u} \begin{pmatrix} 2ue_1 & 0 & \sqrt{2u^3} & 0 & \sqrt{2u^3} \\ 0 & 0 & 0 & 2w & 2\sqrt{v} \\ \sqrt{2u^3} & 0 & 0 & 2\sqrt{v} & 2r \\ 0 & 2w & 2\sqrt{v} & 0 & 0 \\ \sqrt{2u^3} & 2\sqrt{v} & 2r & 0 & 0 \end{pmatrix},$$

$$B(\sigma, g) = \begin{pmatrix} g & \sqrt{l} & \sqrt{m} & 0 & 0 \\ \sqrt{l} & 0 & 0 & 0 & k \\ \sqrt{m} & 0 & e_1 - 2g & \sqrt{m} & 0 \\ 0 & 0 & \sqrt{m} & g & \sqrt{l} \\ 0 & k & 0 & \sqrt{l} & 0 \end{pmatrix},$$

where

$$\begin{aligned} e_n = e_n(\sigma) &= \sum_{1 \leq j_1 < j_2 < \dots < j_n \leq 5} \prod_{i=1}^n \lambda_{j_i} \quad (n = 1, 2, \dots, 5), \\ u = u(\sigma) &= -e_2(\sigma) - \lambda_2^2 - \lambda_5^2, \\ v = v(\sigma) &= -(\lambda_3 + \lambda_5)(\lambda_4 + \lambda_5)(\lambda_2 + \lambda_4)(\lambda_2 + \lambda_3)(\lambda_1 + \lambda_2)(\lambda_1 + \lambda_5), \\ w = w(\sigma) &= \lambda_2 \lambda_5 e_1(\sigma) - \lambda_1 \lambda_3 \lambda_4, \\ r = r(\sigma) &= e_3(\sigma) + e_1(\sigma)(\lambda_2^2 + \lambda_5^2), \\ k = k(\sigma, g) &= g - \lambda_3 - \lambda_5, \\ l = l(\sigma, g) &= (g - \lambda_3)(\lambda_5 - g), \\ m = m(\sigma, g) &= -g^2 + e_1(\sigma)g - \frac{1}{2}(e_2(\sigma) + \lambda_3^2 + \lambda_5^2). \end{aligned}$$

In order for $A(\sigma)$ to be well-defined we assume that $u \neq 0$ and that if $z = r \exp(i\varphi) \in \mathbb{C}$, where r is a nonnegative real number and $-\pi < \varphi \leq \pi$, then $\sqrt{z} = \sqrt{r} \exp(i\frac{\varphi}{2})$.

Lemma 3. Assume $\sigma = (\lambda_1, \lambda_2, \dots, \lambda_5)$ is a list of complex numbers and $u(\sigma) \neq 0$. Then the spectrum of $A(\sigma)$ is σ .

Proof. The characteristic polynomial of $A(\sigma)$ is

$$P_A(z) = z^5 + q_4 z^4 + q_3 z^3 + q_2 z^2 + q_1 z + q_0,$$

where

$$\begin{aligned}
q_4 &= -e_1, \\
q_3 &= -\frac{u^3 + 2v + r^2 + w^2}{u^2}, \\
q_2 &= \frac{-ru^2 + 2e_1v + e_1r^2 + e_1w^2}{u^2}, \\
q_1 &= \frac{-2vwr + w^2u^3 + u^3v + v^2 + r^2w^2}{u^4}, \\
q_0 &= -\frac{-2e_1vwr + u^2vw - u^2w^2r + e_1r^2w^2 + e_1v^2}{u^4}.
\end{aligned}$$

The following expressions are polynomials in $\lambda_1, \dots, \lambda_5$. Therefore, it is straightforward to check that

$$\begin{aligned}
-u^2q_3 &= u^3 + 2v + r^2 + w^2 = -u^2e_2, \\
u^2q_2 &= -ru^2 + 2e_1v + e_1r^2 + e_1w^2 = -u^2e_3, \\
u^4q_1 &= -2vwr + w^2u^3 + u^3v + v^2 + r^2w^2 = u^4e_4, \\
-u^4q_0 &= -2e_1vwr + u^2vw - u^2w^2r + e_1r^2w^2 + e_1v^2 = u^4e_5.
\end{aligned}$$

As $u \neq 0$ we get that

$$\begin{aligned}
P_A(z) &= z^5 - e_1z^4 + e_2z^3 - e_3z^2 + e_4z - e_5 \\
&= (z - \lambda_1)(z - \lambda_2)(z - \lambda_3)(z - \lambda_4)(z - \lambda_5)
\end{aligned}$$

and so the spectrum of $A(\sigma)$ is σ . □

Let

$$Q_\sigma(z) = 2z^3 - 2(\lambda_3 + \lambda_5)z^2 - \left(e_2 + (\lambda_3 - \lambda_5)^2\right)z + e_3 + e_1(\lambda_3^2 + \lambda_5^2).$$

Lemma 4. Assume $\sigma = (\lambda_1, \lambda_2, \dots, \lambda_5)$ is a list of complex numbers and let g be a root of the polynomial $Q_\sigma(z)$. Then the spectrum of $B(\sigma, g)$ is σ .

Proof. The characteristic polynomial of $B(\sigma, g)$ is

$$P_B(z) = z^5 + q_4z^4 + q_3z^3 + q_2z^2 + q_1z + q_0,$$

where

$$\begin{aligned}
q_4 &= -e_1, \\
q_3 &= -k^2 - 2l - 2m + 2ge_1 - 3g^2 = e_2, \\
q_2 &= -2gl + e_1k^2 + 2le_1 + 2mg - g^2e_1 + 2g^3 = Q_\sigma(g) - e_3, \\
q_1 &= 4g^2l + 2k^2m - 2k^2ge_1 + 3k^2g^2 + 2ml - 2gle_1 + l^2 \\
&= -(\lambda_3 + \lambda_5)Q_\sigma(g) + e_4, \\
q_0 &= -2lkm + 2gl^2 - 2k^2mg + k^2g^2e_1 - 2k^2g^3 - l^2e_1 \\
&= \lambda_3\lambda_5Q_\sigma(g) - e_5.
\end{aligned}$$

As we assumed $Q_\sigma(g) = 0$, we get that $q_2 = -e_3$, $q_1 = e_4$ and $q_0 = -e_5$. Therefore,

$$\begin{aligned} P_B(z) &= z^5 - e_1 z^4 + e_2 z^3 - e_3 z^2 + e_4 z - e_5 \\ &= (z - \lambda_1)(z - \lambda_2)(z - \lambda_3)(z - \lambda_4)(z - \lambda_5) \end{aligned}$$

and so the spectrum of $B(\sigma, g)$ is σ . $\square$

4 Unknown Realizable Region

In this section we show what part of the unknown region the new patterns realize.

Lemma 5. *Let $\sigma = (\lambda_1, \lambda_2, \dots, \lambda_5)$ be a list of real numbers. Assume (2) and*

$$\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_5 > -\lambda_1, \quad (5)$$

$$\lambda_3 > e_1(\sigma), \quad (6)$$

$$r(\sigma) \geq 0. \quad (7)$$

Then $A(\sigma)$ is nonnegative, and so σ is realizable by a nonnegative symmetric matrix.

Proof. By (2) element $(1, 1)$ of $A(\sigma)$ is nonnegative. It suffices to show that $u(\sigma) > 0$, $v(\sigma) \geq 0$ and $w(\sigma) \geq 0$ in order to show that $A(\sigma)$ is nonnegative.

We first note that $v(\sigma) > 0$, because by (5), (2) and (6) we have $\lambda_3 > 0 > \lambda_4$ and $\lambda_2 + \lambda_4 < 0$. Therefore, the first three terms of $v(\sigma)$ are negative and the last three terms are positive. Together with the minus sign of $v(\sigma)$ we get a positive value.

By (5), (2) and (6) $\lambda_1 > 0$. Note that

$$u(\sigma) = \lambda_1^2 u\left(\frac{1}{\lambda_1} \cdot \sigma\right), \quad w(\sigma) = \lambda_1^3 w\left(\frac{1}{\lambda_1} \cdot \sigma\right).$$

Therefore, it suffices to show that $u(\sigma) > 0, w(\sigma) \geq 0$ under the assumption that $\lambda_1 = 1$.

We define a different parametrization of the spectrum σ :

$$x = \lambda_2, \quad y = \lambda_3, \quad d = \lambda_2 + \lambda_3 + \lambda_4, \quad t = e_1(\sigma).$$

Then, $\sigma = (1, x, y, d - x - y, -d - 1 + t)$ and $1 \geq x \geq y \geq d - x - y \geq -d - 1 + t > -1$. By (5) $d < t$ and by (6) $y > t$, so $y \in (t, \min(x, -x + 2d + 1 - t)]$.

For given x, d, t we look at u, w as functions of $y \in \mathbb{R}$. Then we have

$$\begin{aligned} u(y) &= y^2 + (x - d)y - xd - d + dt + t - t^2, \\ w(y) &= y^2 + (x - d)y - xdt - tx + t^2x. \end{aligned}$$

Obviously, these functions have an extremum at $y = \frac{1}{2}(d - x)$. By (5) and (6) we get $-x + d - t < 0$, so $u(y)$ and $w(y)$ have a minimum at this point. Moreover, as $u(y)$ and $w(y)$ are quadratic polynomials in y , they are monotonically increasing when $y > \frac{1}{2}(d - x)$.

We have $x \geq y > t > d$, so $\frac{1}{2}(d-x) < 0$. By (2) $t \geq 0$, so $t > \frac{1}{2}(d-x)$. Therefore, if we can show that for $y = t$, $u(y) > 0$ and $w(y) \geq 0$, then it will be true for $y > t$ as well.

In case $y = t$ we get

$$\begin{aligned} u(t) &= -(1+x)(d-t), \\ w(t) &= -t(1+x)(d-t). \end{aligned}$$

As $t \geq 0$, $t > d$ and $x > 0$, indeed $u(t) > 0$ and $w(t) \geq 0$. Therefore, $A(\sigma)$ is nonnegative. By Lemma 3 the spectrum of the symmetric matrix $A(\sigma)$ is σ , so σ is realizable by a nonnegative symmetric matrix. $\square$

The condition $\lambda_5 > -\lambda_1$ in Lemma 5 is crucial, as the following lemma shows:

Lemma 6. *Let $\sigma = (\lambda_1, \lambda_2, \dots, \lambda_5)$ be a list of real numbers. Assume σ meets conditions (2), (6) and*

$$\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_5 = -\lambda_1. \quad (8)$$

Then σ is not realizable by a 5×5 nonnegative matrix.

Proof. By (2), (6) and (8) we have that $\lambda_3 > 0$ and $\lambda_3 + \lambda_4 \leq \lambda_2 + \lambda_4 < 0$. Assume that σ is realizable by a 5×5 nonnegative matrix A . By the Perron-Frobenius Theorem $\lambda_1 + \lambda_4 \geq 0$, and so $\lambda_2 < \lambda_1$. If A is irreducible then it has an index of cyclicity h . As the spectrum of A contains only real values and as $\lambda_5 = -\lambda_1$, then by the Perron-Frobenius Theorem we must have that $h = 2$ and that σ is invariant under multiplication by -1 . However, this contradicts the fact that $\lambda_3 > 0$. If A is reducible then σ can be partitioned into disjoint subsets (in terms of element names and not necessarily element values) forming the spectra of irreducible nonnegative matrices. A subset containing λ_5 must contain λ_1 by the Perron-Frobenius Theorem. Such a subset of σ must also be invariant under multiplication by -1 . As no other element of σ is zero and no two other elements of σ are opposite to one another, such a subset cannot contain any other element of σ . On the other hand, a subset containing λ_4 must contain λ_1 by the Perron-Frobenius Theorem. Therefore, σ cannot be partitioned as required, so A cannot be reducible. We conclude that no 5×5 nonnegative matrix A can realize σ . $\square$

Lemma 7. *Let $\sigma = (\lambda_1, \lambda_2, \dots, \lambda_5)$ be a list of real numbers. Assume σ meets conditions (2), (6) and (4). Further assume that g is a real root of the polynomial $Q_\sigma(z)$ and*

$$0 \leq g \leq \frac{1}{2}e_1(\sigma). \quad (9)$$

Then $B(\sigma, g)$ is nonnegative, and so σ is realizable by a nonnegative symmetric matrix.

Proof. By (9) the main diagonal of $B(\sigma, g)$ is nonnegative. By (4), (2) and (6) we have $\lambda_3 > e_1(\sigma) \geq \frac{1}{2}e_1(\sigma) \geq g \geq 0 > \lambda_5$ and $\lambda_3 + \lambda_5 < 0 \leq g$. Therefore,

$l(\sigma, g) > 0$ and $k(\sigma, g) > 0$. Note that $m(\sigma, z)$ can be regarded as a quadratic polynomial in z . Its discriminant is

$$\begin{aligned}\Delta &= e_1(\sigma)^2 - 2(e_2(\sigma) + \lambda_3^2 + \lambda_5^2) \\ &= (\lambda_4^2 - \lambda_3^2) + \lambda_2^2 + (\lambda_1^2 - \lambda_5^2).\end{aligned}$$

By (4), (2) and (6) we have $-\lambda_1 \leq \lambda_5 \leq \lambda_4 < -\lambda_3 < 0$. Therefore, $\Delta \geq \lambda_2^2 > 0$ and so $m(\sigma, z)$ has two real roots:

$$z_0^\pm = \frac{1}{2}e_1(\sigma) \pm \frac{1}{2}\sqrt{\Delta}.$$

As the coefficient of z^2 in $m(\sigma, z)$ is negative, $m(\sigma, z) \geq 0$ for any $z \in [z_0^-, z_0^+]$. Also, we know that

$$\begin{aligned}z_0^- &\leq \frac{1}{2}e_1(\sigma) - \frac{1}{2}\lambda_2 \\ &= \frac{1}{2}(\lambda_1 + \lambda_3 + \lambda_4 + \lambda_5) \\ &\leq \frac{1}{2}(\lambda_1 + \lambda_2 + \lambda_4 + \lambda_5) < 0 \\ &\leq g \leq \frac{1}{2}e_1(\sigma) \leq z_0^+.\end{aligned}$$

Therefore, $m(\sigma, g) \geq 0$, and so $B(\sigma, g)$ is nonnegative. By Lemma 4 the spectrum of the symmetric matrix $B(\sigma, g)$ is σ , so σ is realizable by a nonnegative symmetric matrix. $\square$

5 Guo Perturbations of Symmetric Nonnegative Matrices

Let the list of complex numbers $\sigma = (\lambda_1, \lambda_2, \dots, \lambda_n)$ be the spectrum of an $n \times n$ nonnegative real matrix with λ_1 its Perron eigenvalue. Let s be a positive real number and let $i \in \{2, 3, \dots, n\}$. We define the following lists of values:

- σ_i^{+s} is the list σ with λ_1 replaced by $\lambda_1 + s$ and λ_i replaced by $\lambda_i + s$,
- σ_i^{-s} is the list σ with λ_1 replaced by $\lambda_1 + s$ and λ_i replaced by $\lambda_i - s$.

In [4] Guo proved that if a list of complex numbers $\sigma = (\lambda_1, \lambda_2, \dots, \lambda_n)$ is the spectrum of an $n \times n$ nonnegative real matrix with λ_1 its Perron eigenvalue and $\lambda_i \in \mathbb{R}$, then σ_i^{+s} and σ_i^{-s} are also realizable by an $n \times n$ nonnegative real matrix. In this section we prove similar results for several families of 5×5 symmetric nonnegative matrices.

In [11] the eigenvalues of trace zero 5×5 symmetric nonnegative matrices were fully characterized. A descending list of real numbers $\lambda_1 \geq \lambda_2 \geq \lambda_3 \geq \lambda_4 \geq \lambda_5 \geq -\lambda_1$ is realizable by a trace zero 5×5 symmetric nonnegative matrix if and only if the following conditions hold:

1. $\sum_{i=1}^5 \lambda_i = 0$,
2. $\sum_{i=1}^5 \lambda_i^3 \geq 0$,

$$3. \lambda_2 + \lambda_5 \leq 0.$$

Theorem 3. *Let a list of real numbers $\sigma = (\lambda_1, \lambda_2, \dots, \lambda_5)$, where $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_5 \geq -\lambda_1$, be the spectrum of a trace zero 5×5 symmetric nonnegative matrix. Then σ_i^{-s} is realizable by a trace zero 5×5 symmetric nonnegative matrix for any positive real number s and any $i \in \{2, 3, 4, 5\}$.*

Proof. The sum of elements of σ and of σ_i^{-s} is the same, so the first condition is met for σ_i^{-s} . As $s > 0$ we know that λ_2 and λ_5 can only get smaller, so the third condition holds for σ_i^{-s} as well. Finally, as $-\lambda_1 \leq \lambda_i \leq \lambda_1$ we get that $\lambda_1 + \lambda_i \geq 0$ and $\lambda_1 - \lambda_i \geq 0$. Therefore,

$$(\lambda_1 + s)^3 + (\lambda_i - s)^3 = 3s(\lambda_1 + \lambda_i)(\lambda_1 + s - \lambda_i) + \lambda_1^3 + \lambda_i^3 \geq \lambda_1^3 + \lambda_i^3.$$

We deduce that the second condition is also met for σ_i^{-s} . Therefore, σ_i^{-s} is realizable by a trace zero 5×5 symmetric nonnegative matrix. $\square$

Theorem 4. *Let a list of real numbers $\sigma = (\lambda_1, \lambda_2, \dots, \lambda_5)$, where $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_5 \geq -\lambda_1$ and $\lambda_3 \leq \sum_{i=1}^5 \lambda_i$, be the spectrum of a 5×5 symmetric nonnegative matrix. Then σ_i^{+s} and σ_i^{-s} are realizable by a 5×5 symmetric nonnegative matrix for any positive real number s and any $i \in \{2, 3, 4, 5\}$.*

Proof. Let $\sigma = (\lambda_1, \lambda_2, \dots, \lambda_5)$ be the spectrum of a 5×5 symmetric nonnegative matrix and meet conditions (4) and

$$\lambda_3 \leq e_1(\sigma). \quad (10)$$

By Lemma 1 condition (3) is met. By (4) we have $\max_{\lambda \in \sigma_i^{+s}} |\lambda| = \lambda_1 + s \in \sigma_i^{+s}$ and $\max_{\lambda \in \sigma_i^{-s}} |\lambda| = \lambda_1 + s \in \sigma_i^{-s}$ for any $i \in \{2, 3, 4, 5\}$. We know that (2) is met for σ , so we have $e_1(\sigma_i^{+s}) \geq e_1(\sigma) \geq 0$ and $e_1(\sigma_i^{-s}) = e_1(\sigma) \geq 0$ for any $i \in \{2, 3, 4, 5\}$. Therefore, conditions (1) and (2) are met by σ_i^{+s} and σ_i^{-s} . We look into the following cases:

$$1. \lambda_3 > 0.$$

$$(a) i \in \{2, 4, 5\}.$$

In this case let σ' be the perturbed σ (σ_i^{+s} or σ_i^{-s}) without the element λ_3 . By (10) $e_1(\sigma') \geq \lambda_1 + \lambda_2 + \lambda_4 + \lambda_5 \geq 0$. Also, $\max_{\lambda \in \sigma'} |\lambda| = \lambda_1 + s \in \sigma'$. Therefore, σ' meets conditions (1) and (2), so by Theorem 2 there exists a 4×4 symmetric nonnegative matrix A_1 realizing σ' . Hence, σ is realizable by the 5×5 symmetric nonnegative matrix $A = A_1 \oplus (\lambda_3)$.

$$(b) i = 3.$$

$$i. \sigma_3^{+s}.$$

This is similar to case 1a with $\sigma' = (\lambda_1 + s, \lambda_2, \lambda_4, \lambda_5)$ and $A = A_1 \oplus (\lambda_3 + s)$.

$$ii. \sigma_3^{-s} \text{ and } \lambda_3 - s \geq 0.$$

This is similar to case 1a with $\sigma' = (\lambda_1 + s, \lambda_2, \lambda_4, \lambda_5)$ and $A = A_1 \oplus (\lambda_3 - s)$.

- iii. σ_3^{-s} and $\lambda_3 - s < 0$ and $\lambda_4 \leq 0$.
 In this case σ_3^{-s} has exactly two positive elements: $\lambda_1 + s$ and λ_2 . If $\lambda_3 - s \geq \lambda_5$ then the sum of the first, third and fourth largest elements in σ_3^{-s} is $(\lambda_1 + s) + (\lambda_3 - s) + \lambda_4 = \lambda_1 + \lambda_3 + \lambda_4$. If $\lambda_3 - s < \lambda_5$ then the sum of the first, third and fourth largest elements in σ_3^{-s} is $(\lambda_1 + s) + \lambda_4 + \lambda_5 > (\lambda_1 + s) + \lambda_4 + (\lambda_3 - s) = \lambda_1 + \lambda_3 + \lambda_4$. In both cases by (3) this sum is nonnegative, so σ_3^{-s} meets condition (3). Therefore, by Lemma 2, σ_3^{-s} is realizable by a 5×5 symmetric nonnegative matrix.
- iv. σ_3^{-s} and $\lambda_3 - s < 0$ and $\lambda_4 > 0$.
 This is similar to case 1a with $\sigma' = (\lambda_1 + s, \lambda_2, \lambda_3 - s, \lambda_5)$ and $A = A_1 \oplus (\lambda_4)$.

2. $\lambda_3 \leq 0$.

(a) $i = 2$.

- i. σ_2^{+s} and $\lambda_2 + s > 0$.
 In this case σ_2^{+s} has exactly two positive elements: $\lambda_1 + s$ and $\lambda_2 + s$. By (3) the sum of the first, third and fourth largest elements in σ_2^{+s} is $(\lambda_1 + s) + \lambda_3 + \lambda_4 > \lambda_1 + \lambda_3 + \lambda_4 \geq 0$, so σ_2^{+s} meets condition (3). Hence, by Lemma 2, σ_2^{+s} is realizable by a 5×5 symmetric nonnegative matrix.
- ii. σ_2^{+s} and $\lambda_2 + s \leq 0$.
 In this case σ_2^{+s} is realizable by Theorem 1, as it meets the conditions of this theorem.
- iii. σ_2^{-s} and $\lambda_2 - s > 0$.
 This is similar to case 2(a)i.
- iv. σ_2^{-s} and $\lambda_2 - s \leq 0$.
 This is similar to case 2(a)ii.

(b) $i \in \{3, 4\}$.

- i. σ_i^{+s} and $\lambda_2 \leq 0$ and $\lambda_i + s \leq 0$.
 This is similar to case 2(a)ii.
- ii. σ_i^{+s} and $\lambda_2 > 0$ and $\lambda_i + s \leq 0$.
 This is similar to case 2(a)i.
- iii. σ_i^{+s} and $\lambda_i + s > 0$.
 This is similar to case 1a with $\sigma' = (\lambda_1 + s, \lambda_2, \lambda_{7-i}, \lambda_5)$ and $A = A_1 \oplus (\lambda_i + s)$.
- iv. σ_i^{-s} and $\lambda_2 > 0$.
 This is similar to case 1(b)iii.
- v. σ_i^{-s} and $\lambda_2 \leq 0$.
 This is similar to case 2(a)ii.

(c) $i = 5$.

- i. σ_5^{+s} and $\lambda_2 \leq 0$ and $\lambda_5 + s \leq 0$.
 This is similar to case 2(a)ii.
- ii. σ_5^{+s} and $\lambda_2 > 0$ and $\lambda_5 + s \leq 0$.
 This is similar to case 1(b)iii.

- iii. σ_5^{+s} and $\lambda_5 + s > 0$ and $\lambda_2 \geq \lambda_5 + s$.
This is similar to case 1a with $\sigma' = (\lambda_1 + s, \lambda_2, \lambda_3, \lambda_4)$ and $A = A_1 \oplus (\lambda_5 + s)$.
- iv. σ_5^{+s} and $\lambda_5 + s > 0$ and $\lambda_2 < \lambda_5 + s$.
This is similar to case 1a with $\sigma' = (\lambda_1 + s, \lambda_5 + s, \lambda_3, \lambda_4)$ and $A = A_1 \oplus (\lambda_2)$.
- v. σ_5^{-s} and $\lambda_2 > 0$.
This is similar to case 2(a)i.
- vi. σ_5^{-s} and $\lambda_2 \leq 0$.
This is similar to case 2(a)ii.

□

Theorem 5. *Let $\sigma = (\lambda_1, \lambda_2, \dots, \lambda_5)$ be a list of real numbers. Assume σ meets the conditions of Lemma 5. Then σ_i^{-s} meets the conditions of Lemma 5 for sufficiently small positive real number s and any $i \in \{2, 3, 4, 5\}$, and so is realizable by a 5×5 symmetric nonnegative matrix.*

Proof. As we wish to prove the result for sufficiently small values of s , we may assume that if $\lambda_i > \lambda_{i+1}$ then $\lambda_i - s \geq \lambda_{i+1}$.

Note that $e_1(\sigma_i^{-s}) = e_1(\sigma) \geq 0$ for any positive real number s and any $i \in \{2, 3, 4, 5\}$, so (2) is valid for σ_i^{-s} .

We divide the proof into several cases:

1. $i = 2$.

(a) $\lambda_2 > \lambda_3$.

We have $\lambda_1 + s > \lambda_2 - s \geq \lambda_3$ and $\lambda_5 > -\lambda_1 > -(\lambda_1 + s)$, so (5) is valid for σ_2^{-s} .

Since λ_3 is the third element of σ_2^{-s} and $\lambda_3 > e_1(\sigma) = e_1(\sigma_2^{-s})$ then (6) is valid for σ_2^{-s} .

If $r(\sigma) > 0$ then, by continuity of $r(\sigma)$, we have $r(\sigma_2^{-s}) > 0$ for sufficiently small s . This means (7) is valid for σ_2^{-s} . Assume $r(\sigma) = 0$. Then

$$r(\sigma_2^{-s}) = r(\sigma_2^{-s}) - r(\sigma) = s(\lambda_1 + \lambda_2)(-2\lambda_2 - \lambda_3 - \lambda_4 - \lambda_5 + s).$$

As $s > 0$ and $\lambda_1 + \lambda_2 > 0$ it suffices to prove that $2\lambda_2 + \lambda_3 + \lambda_4 + \lambda_5 \leq 0$ in order to have $r(\sigma_2^{-s}) > 0$. This will show (7) is valid for σ_2^{-s} .

Collecting terms we get

$$\begin{aligned} r(\sigma) &= r_4(\sigma)\lambda_4 + \lambda_1\lambda_2\lambda_3 + \lambda_1\lambda_3\lambda_5 + \lambda_2\lambda_3\lambda_5 + \lambda_1\lambda_2\lambda_5 \\ &\quad + (\lambda_1 + \lambda_2 + \lambda_3 + \lambda_5)(\lambda_2^2 + \lambda_5^2), \end{aligned}$$

where

$$r_4(\sigma) = (\lambda_2 + \lambda_3 + \lambda_5)\lambda_1 + \lambda_3\lambda_5 + \lambda_2\lambda_3 + \lambda_2\lambda_5 + \lambda_5^2 + \lambda_2^2.$$

Assume $r_4(\sigma) = 0$. If $\lambda_2 + \lambda_3 + \lambda_5 = 0$ then

$$\lambda_3\lambda_5 + \lambda_2\lambda_3 + \lambda_2\lambda_5 + \lambda_5^2 + \lambda_2^2 = 0. \quad (11)$$

Substituting $\lambda_5 = -\lambda_2 - \lambda_3$ in (11) we get $\lambda_2(\lambda_2 + \lambda_3) = 0$, which is impossible because by (6), $\lambda_2 \geq \lambda_3 > 0$. Therefore, $\lambda_2 + \lambda_3 + \lambda_5 \neq 0$ and so,

$$\lambda_1 = -\frac{\lambda_3\lambda_5 + \lambda_2\lambda_3 + \lambda_2\lambda_5 + \lambda_5^2 + \lambda_2^2}{\lambda_2 + \lambda_3 + \lambda_5}. \quad (12)$$

Substituting (12) in $u(\sigma)$ gives

$$u(\sigma) = -\frac{\lambda_4(\lambda_3 + \lambda_5)(\lambda_2 + \lambda_3)}{\lambda_2 + \lambda_3 + \lambda_5}.$$

By (6), $\lambda_4 < 0$, $\lambda_2 + \lambda_3 > 0$ and $\lambda_3 + \lambda_5 < 0$. By Lemma 5, $u(\sigma) > 0$. Therefore, we must have $\lambda_2 + \lambda_3 + \lambda_5 < 0$.

Assume $r_4(\sigma) \neq 0$. Then, as we assumed $r(\sigma) = 0$,

$$\begin{aligned} \lambda_4 = & -\frac{\lambda_1\lambda_2\lambda_3 + \lambda_1\lambda_3\lambda_5 + \lambda_2\lambda_3\lambda_5 + \lambda_1\lambda_2\lambda_5}{r_4(\sigma)} \\ & -\frac{(\lambda_1 + \lambda_2 + \lambda_3 + \lambda_5)(\lambda_2^2 + \lambda_5^2)}{r_4(\sigma)}. \end{aligned} \quad (13)$$

Substituting (13) in $u(\sigma)$ gives

$$u(\sigma) = -\frac{(\lambda_3 + \lambda_5)(\lambda_2 + \lambda_3)(\lambda_1 + \lambda_5)(\lambda_1 + \lambda_2)}{r_4(\sigma)}. \quad (14)$$

By (5) and (6), the left term in the numerator of (14) is negative and the rest of the terms in the numerator are positive. By Lemma 5, $u(\sigma) > 0$. Therefore, we must have that $r_4(\sigma) > 0$. Also, by (6), $\lambda_1 + \lambda_2 + \lambda_4 + \lambda_5 < 0$. Substituting (13) in this expression we get

$$\frac{(\lambda_1 + \lambda_5)(\lambda_1 + \lambda_2)(\lambda_2 + \lambda_3 + \lambda_5)}{r_4(\sigma)} < 0. \quad (15)$$

As we already know that $r_4(\sigma) > 0$ and that the first two terms of the numerator of (15) are positive, then we must have $\lambda_2 + \lambda_3 + \lambda_5 < 0$. Regardless of the value of $r_4(\sigma)$ we have shown that $\lambda_2 + \lambda_3 + \lambda_5 < 0$. Therefore, $2\lambda_2 + \lambda_3 + \lambda_4 + \lambda_5 < \lambda_2 + \lambda_4 < 0$.

(b) $\lambda_2 = \lambda_3 > \lambda_4$.

This case is covered by case 2a.

(c) $\lambda_2 = \lambda_3 = \lambda_4 > \lambda_5$.

This case is covered by case 3a.

(d) $\lambda_2 = \lambda_3 = \lambda_4 = \lambda_5$.

This case is covered by case 4.

2. $i = 3$.

(a) $\lambda_3 > \lambda_4$.

We have $\lambda_1 + s > \lambda_2 > \lambda_3 - s \geq \lambda_4$ and $\lambda_5 > -\lambda_1 > -(\lambda_1 + s)$, so (5) is valid for σ_3^{-s} .

Since $\lambda_3 - s$ is the third element of σ_3^{-s} and $\lambda_3 > e_1(\sigma) = e_1(\sigma_3^{-s})$ we require that $s < \lambda_3 - e_1(\sigma)$, so that (6) is valid for σ_3^{-s} .

We also have

$$r(\sigma_3^{-s}) - r(\sigma) = -s(\lambda_2 + \lambda_4 + \lambda_5)(s + \lambda_1 - \lambda_3).$$

We know $s > 0$. By (5) $\lambda_1 \geq \lambda_3$ and by (6) $\lambda_2 + \lambda_4 + \lambda_5 < 0$, so $r(\sigma_3^{-s}) - r(\sigma) > 0$. This shows (7) is valid for σ_3^{-s} .

(b) $\lambda_3 = \lambda_4 > \lambda_5$.

This case is covered by case 3a.

(c) $\lambda_3 = \lambda_4 = \lambda_5$.

This case is covered by case 4.

3. $i = 4$.

(a) $\lambda_4 > \lambda_5$.

We have $\lambda_1 + s > \lambda_2$, $\lambda_4 - s \geq \lambda_5$ and $\lambda_5 > -\lambda_1 > -(\lambda_1 + s)$, so (5) is valid for σ_4^{-s} .

Since λ_3 is the third element of σ_4^{-s} and $\lambda_3 > e_1(\sigma) = e_1(\sigma_4^{-s})$ then (6) is valid for σ_4^{-s} .

If $r(\sigma) > 0$ then, by continuity of $r(\sigma)$, we have $r(\sigma_4^{-s}) > 0$ for sufficiently small s . This means (7) is valid for σ_4^{-s} . Assume $r(\sigma) = 0$. Then

$$r(\sigma_4^{-s}) - r(\sigma) = -s(\lambda_2 + \lambda_3 + \lambda_5)(s + \lambda_1 - \lambda_4).$$

We know $s > 0$ and by (5) $s + \lambda_1 - \lambda_4 > 0$. In case 1a we already showed that when $r(\sigma) = 0$ we have $\lambda_2 + \lambda_3 + \lambda_5 < 0$. Therefore, $r(\sigma_4^{-s}) > 0$. This shows (7) is valid for σ_4^{-s} .

(b) $\lambda_4 = \lambda_5$.

This case is covered by case 4.

4. $i = 5$.

We have $\lambda_1 + s > \lambda_2$, and by (5) $\lambda_5 - s > -(\lambda_1 + s)$, so (5) is valid for σ_5^{-s} .

Since λ_3 is the third element of σ_5^{-s} and $\lambda_3 > e_1(\sigma) = e_1(\sigma_5^{-s})$ then (6) is valid for σ_5^{-s} .

We also have

$$r(\sigma_5^{-s}) - r(\sigma) = s(\lambda_1 + \lambda_5)(s - \lambda_2 - \lambda_3 - \lambda_4 - 2\lambda_5).$$

We know $s > 0$. By (5) $\lambda_1 + \lambda_5 > 0$ and by (6) $\lambda_2 + \lambda_3 + \lambda_4 + 2\lambda_5 < 0$, so $r(\sigma_5^{-s}) - r(\sigma) > 0$. This shows (7) is valid for σ_5^{-s} .

□

Theorem 6. Let $\sigma = (\lambda_1, \lambda_2, \dots, \lambda_5)$ be a list of real numbers and $g \in \mathbb{R}$. Assume σ and g meet the conditions of Lemma 7 and also the condition

$$r(\sigma) < 0. \tag{16}$$

Then, for sufficiently small positive real number s and $i \in \{2, 3, 4, 5\}$, there exists $\tilde{g} \in \mathbb{R}$ (depending on s and i) such that σ_i^{-s} and $\tilde{g}$ meet the conditions of Lemma 7 and (16), and so σ_i^{-s} is realizable by a 5×5 symmetric nonnegative matrix.

Proof. As we wish to prove the result for sufficiently small values of s , we may assume that if $\lambda_i > \lambda_{i+1}$ then $\lambda_i - s \geq \lambda_{i+1}$.

Note that $e_1(\sigma_i^{-s}) = e_1(\sigma) \geq 0$ for any positive real number s and any $i \in \{2, 3, 4, 5\}$, so (2) is valid for σ_i^{-s} .

By our assumption $r(\sigma) < 0$. Also, by continuity of $r(\sigma)$ in λ_1 and λ_i , for sufficiently small positive real number s we have $r(\sigma_i^{-s}) < 0$. Therefore, (16) is valid for σ_i^{-s} .

By our assumptions there is $g \in [0, \frac{1}{2}e_1(\sigma)]$ such that $Q_\sigma(g) = 0$ and $B(\sigma, g)$ is nonnegative. By (4), (2) and (6) we know that $\lambda_2 \geq \lambda_3 > 0$. Therefore, by (16) we have

$$e_3(\sigma) + e_1(\sigma) (\lambda_3^2 + \lambda_5^2) \leq e_3(\sigma) + e_1(\sigma) (\lambda_2^2 + \lambda_5^2) = r(\sigma) < 0,$$

which means that the $Q_\sigma(0) < 0$. By continuity of polynomials, for sufficiently small positive real number s and $i \in \{2, 3, 4, 5\}$, $Q_{\sigma_i^{-s}}(0) < 0$.

The inflection point of $Q_\sigma(z)$ is at $\frac{1}{3}(\lambda_3 + \lambda_5)$. By (4), (2) and (6), $\lambda_3 + \lambda_5 < 0$ and as the coefficient of z^3 in $Q_\sigma(z)$ is positive, then for $z \geq 0$ the polynomial $Q_\sigma(z)$ is convex. This means that there can only be a single real root of $Q_\sigma(z)$ in the range $(0, \frac{1}{2}e_1(\sigma)]$.

If $g < \frac{1}{2}e_1(\sigma)$ then, by continuity of the roots of a polynomial, for sufficiently small positive real number s and $i \in \{2, 3, 4, 5\}$, $Q_{\sigma_i^{-s}}(z)$ has a real root $\tilde{g} \in (0, \frac{1}{2}e_1(\sigma))$ so (9) is valid for $Q_{\sigma_i^{-s}}(z)$. We will therefore assume that

$$Q_\sigma\left(\frac{1}{2}e_1(\sigma)\right) = 0. \quad (17)$$

We divide the proof into several cases:

1. $i = 2$.

(a) $\lambda_2 > \lambda_3$.

We have $\lambda_1 + s > \lambda_2 - s \geq \lambda_3$ and $\lambda_5 \geq -\lambda_1 > -(\lambda_1 + s)$, so (4) is valid for σ_2^{-s} .

Since λ_3 is the third element of σ_2^{-s} and $\lambda_3 > e_1(\sigma) = e_1(\sigma_2^{-s})$ then (6) is valid for σ_2^{-s} .

We have

$$Q_{\sigma_2^{-s}}(z) - Q_\sigma(z) = s(s + \lambda_1 - \lambda_2)(z - \lambda_3 - \lambda_4 - \lambda_5).$$

By (4), (2) and (6), $\lambda_1 \geq \lambda_2$ and $\lambda_3 + \lambda_4 + \lambda_5 < 0$. Together with $s > 0$ we get that the above expression is positive for $z \geq 0$, and in particular for $z = \frac{1}{2}e_1(\sigma)$, so that by (17)

$$Q_{\sigma_2^{-s}}\left(\frac{1}{2}e_1(\sigma)\right) = Q_{\sigma_2^{-s}}\left(\frac{1}{2}e_1(\sigma)\right) - Q_\sigma\left(\frac{1}{2}e_1(\sigma)\right) > 0.$$

Thus, $Q_{\sigma_2^{-s}}(z)$ must have a real root $\tilde{g} \in (0, \frac{1}{2}e_1(\sigma))$, so (9) is valid for $Q_{\sigma_2^{-s}}(z)$.

(b) $\lambda_2 = \lambda_3 > \lambda_4$.

This case is covered by case 2a.

(c) $\lambda_2 = \lambda_3 = \lambda_4 > \lambda_5$.

This case is covered by case 3a.

(d) $\lambda_2 = \lambda_3 = \lambda_4 = \lambda_5$.

This case is covered by case 4.

2. $i = 3$.

(a) $\lambda_3 > \lambda_4$.

We have $\lambda_1 + s > \lambda_2 > \lambda_3 - s \geq \lambda_4$ and $\lambda_5 \geq -\lambda_1 > -(\lambda_1 + s)$, so (4) is valid for σ_3^{-s} .

Since $\lambda_3 - s$ is the third element of σ_3^{-s} and $\lambda_3 > e_1(\sigma) = e_1(\sigma_3^{-s})$ we require that $s < \lambda_3 - e_1(\sigma)$, so that (6) is valid for σ_3^{-s} .

We have

$$\begin{aligned} Q_{\sigma_3^{-s}} \left(\frac{1}{2} e_1(\sigma) \right) - Q_{\sigma} \left(\frac{1}{2} e_1(\sigma) \right) &= (\lambda_1 + \lambda_3) s^2 \\ &+ \frac{1}{2} s \left((\lambda_2 - \lambda_3 + \lambda_4 - \lambda_5) (\lambda_1 + \lambda_3) + (\lambda_2 + \lambda_4)^2 \right) \\ &+ \frac{1}{2} s \left((\lambda_1 - \lambda_5) (\lambda_1 + \lambda_5) + \lambda_1^2 + \lambda_3 (\lambda_1 - \lambda_3) \right). \end{aligned}$$

By (4) we have $\lambda_1 \geq \lambda_2 \geq \lambda_3 \geq \lambda_4 \geq \lambda_5 \geq -\lambda_1$, and by (2) and (6) $\lambda_1 \geq \lambda_3 > 0$. Together with $s > 0$ we get that the above expression is positive. Hence, by (17)

$$Q_{\sigma_3^{-s}} \left(\frac{1}{2} e_1(\sigma) \right) = Q_{\sigma_3^{-s}} \left(\frac{1}{2} e_1(\sigma) \right) - Q_{\sigma} \left(\frac{1}{2} e_1(\sigma) \right) > 0,$$

and so $Q_{\sigma_3^{-s}}(z)$ must have a real root $\tilde{g} \in (0, \frac{1}{2} e_1(\sigma))$. Therefore, (9) is valid for $Q_{\sigma_3^{-s}}(z)$.

(b) $\lambda_3 = \lambda_4 > \lambda_5$.

This case is covered by case 3a.

(c) $\lambda_3 = \lambda_4 = \lambda_5$.

This case is covered by case 4.

3. $i = 4$.

(a) $\lambda_4 > \lambda_5$.

We have $\lambda_1 + s > \lambda_2$, $\lambda_4 - s \geq \lambda_5$ and $\lambda_5 \geq -\lambda_1 > -(\lambda_1 + s)$, so (4) is valid for σ_4^{-s} .

Since λ_3 is the third element of σ_4^{-s} and $\lambda_3 > e_1(\sigma) = e_1(\sigma_4^{-s})$ then (6) is valid for σ_4^{-s} .

Note that

$$\begin{aligned} Q_{\sigma} \left(\frac{1}{2} e_1(\sigma) \right) &= \frac{1}{2} \lambda_2 (\lambda_2 + \lambda_5) (\lambda_1 + \lambda_2 + \lambda_3 + \lambda_4 - \lambda_5) \\ &- \frac{1}{2} (\lambda_1 - \lambda_2 - \lambda_3 + \lambda_4 - \lambda_5) (\lambda_1 + \lambda_3) (\lambda_3 + \lambda_4) \\ &+ \frac{1}{4} (\lambda_1 - \lambda_2 - \lambda_3 + \lambda_4 - \lambda_5) (\lambda_1 + \lambda_2 + \lambda_3 + \lambda_4 - \lambda_5) e_1(\sigma). \end{aligned}$$

If $\lambda_1 - \lambda_2 - \lambda_3 + \lambda_4 - \lambda_5 \leq 0$, then together with (4), (2) and (6) we get that the above expression is negative. This contradicts (17), and so

$$\lambda_1 - \lambda_2 - \lambda_3 + \lambda_4 - \lambda_5 > 0. \quad (18)$$

We have

$$\begin{aligned} Q_{\sigma_4^{-s}} \left(\frac{1}{2} e_1(\sigma) \right) - Q_\sigma \left(\frac{1}{2} e_1(\sigma) \right) = \\ \frac{1}{2} s (\lambda_1 - \lambda_2 - \lambda_3 + \lambda_4 - \lambda_5) (s + \lambda_1 - \lambda_4). \end{aligned}$$

By (4), (17), (18) and the fact that $s > 0$ we have

$$Q_{\sigma_4^{-s}} \left(\frac{1}{2} e_1(\sigma) \right) = Q_{\sigma_4^{-s}} \left(\frac{1}{2} e_1(\sigma) \right) - Q_\sigma \left(\frac{1}{2} e_1(\sigma) \right) > 0.$$

Thus, $Q_{\sigma_4^{-s}}(z)$ must have a real root $\tilde{g} \in (0, \frac{1}{2} e_1(\sigma))$, so (9) is valid for $Q_{\sigma_4^{-s}}(z)$.

(b) $\lambda_4 = \lambda_5$.

This case is covered by case 4.

4. $i = 5$.

We have $\lambda_1 + s > \lambda_2$, and by (5) $\lambda_5 - s \geq -(\lambda_1 + s)$, so (4) is valid for σ_5^{-s} .

Since λ_3 is the third element of σ_5^{-s} and $\lambda_3 > e_1(\sigma) = e_1(\sigma_5^{-s})$ then (6) is valid for σ_5^{-s} .

Note that

$$\begin{aligned} Q_\sigma \left(\frac{1}{2} e_1(\sigma) \right) = \\ \lambda_1 \lambda_2 \lambda_4 + \frac{1}{4} (\lambda_1 + \lambda_2 - \lambda_3 + \lambda_4 - \lambda_5) (\lambda_1^2 + \lambda_2^2 - \lambda_3^2 + \lambda_4^2 - \lambda_5^2). \end{aligned}$$

By (4), (2) and (6) $\lambda_1 + \lambda_2 - \lambda_3 + \lambda_4 - \lambda_5 > 0$, so by (17) we get

$$\lambda_1^2 + \lambda_2^2 - \lambda_3^2 + \lambda_4^2 - \lambda_5^2 = -\frac{4\lambda_1 \lambda_2 \lambda_4}{\lambda_1 + \lambda_2 - \lambda_3 + \lambda_4 - \lambda_5}.$$

Therefore, by (4), (2) and (6)

$$\begin{aligned} \lambda_2 \lambda_4 + \frac{1}{2} (\lambda_1^2 + \lambda_2^2 - \lambda_3^2 + \lambda_4^2 - \lambda_5^2) = \\ -\frac{\lambda_2 \lambda_4 (\lambda_3 - \lambda_2 - \lambda_4 + \lambda_1 + \lambda_5)}{\lambda_1 + \lambda_2 - \lambda_3 + \lambda_4 - \lambda_5} > 0. \end{aligned} \quad (19)$$

We have

$$\begin{aligned} Q_{\sigma_5^{-s}} \left(\frac{1}{2} e_1(\sigma) \right) - Q_\sigma \left(\frac{1}{2} e_1(\sigma) \right) = (\lambda_1 + \lambda_5) s^2 \\ + \frac{1}{2} s (\lambda_1 + \lambda_5) (\lambda_1 + \lambda_2 - \lambda_3 + \lambda_4 - \lambda_5) \\ + s \left(\lambda_2 \lambda_4 + \frac{1}{2} (\lambda_1^2 + \lambda_2^2 - \lambda_3^2 + \lambda_4^2 - \lambda_5^2) \right). \end{aligned}$$

By (4), (2), (6), (17), (19) and the fact that $s > 0$ we have

$$Q_{\sigma_5^{-s}}\left(\frac{1}{2}e_1(\sigma)\right) = Q_{\sigma_5^{-s}}\left(\frac{1}{2}e_1(\sigma)\right) - Q_{\sigma}\left(\frac{1}{2}e_1(\sigma)\right) > 0.$$

Thus, $Q_{\sigma_5^{-s}}(z)$ must have a real root $\tilde{g} \in (0, \frac{1}{2}e_1(\sigma))$, so (9) is valid for $Q_{\sigma_5^{-s}}(z)$.

□

Lemma 8. *Let A be a symmetric nonnegative matrix with a spectral radius $\rho(A)$. Then the elements of A are in the range $[0, \rho(A)]$.*

Proof. It is well-known that for an $n \times n$ symmetric matrix $A = (a_{ij})$ we have

$$\rho(A) = \max_{\|x\|_2=1} \|Ax\|_2.$$

Let $\mathbf{e}_i$ be the i^{th} unit vector in $\mathbb{R}^n$. Then, as A is nonnegative,

$$0 \leq a_{ij} = \sqrt{a_{ij}^2} \leq \sqrt{\sum_{k=1}^n a_{kj}^2} = \|A\mathbf{e}_j\|_2 \leq \rho(A).$$

□

In the next theorem we do not assume that the elements of σ are monotonically decreasing.

Theorem 7. *Let $i \in \{2, 3, \dots, n\}$ be fixed. Assume that if a list of real numbers $\sigma = (\lambda_1, \lambda_2, \dots, \lambda_n)$ is realizable by a nonnegative symmetric matrix with Perron eigenvalue λ_1 , then there exists a positive real number s_0 (depending on σ and i), such that σ_i^{+s} (σ_i^{-s}) is realizable by a nonnegative symmetric matrix with Perron eigenvalue $\lambda_1 + s$ for all $0 \leq s < s_0$. Then σ_i^{+s} (σ_i^{-s}) is realizable by a symmetric nonnegative matrix for any positive real number s .*

Proof. We follow the path of Laffey in [6]. Let $i \in \{2, 3, \dots, n\}$ be fixed and let $\sigma = (\lambda_1, \lambda_2, \dots, \lambda_n)$ be a fixed realizable spectrum of a nonnegative symmetric matrix. Assume that for some $r > 0$, σ_i^{+r} (σ_i^{-r}) is not realizable by a symmetric nonnegative matrix. Let

$$W^{\pm} = \{t > 0 \mid \sigma_i^{\pm s} \text{ is realizable for all } s \text{ with } 0 \leq s < t\}.$$

Note that by assumption, there exists a positive real number s_0 such that $s_0 \in W^+$ ($s_0 \in W^-$). Also, W^+ (W^-) is bounded above by r , so it has a supremum, which we denote by w . Obviously, $w \geq s_0 > 0$. Let $\{w_k\}$ ($k = 1, 2, \dots$) be a strictly increasing sequence of positive real numbers with limit w . Note that $\sigma_i^{+w_k}$ ($\sigma_i^{-w_k}$) is realizable by a symmetric nonnegative matrix, or else $w \leq w_k$, in contradiction of the definition of the sequence $\{w_k\}$. Let A_k be a symmetric nonnegative matrix having the spectrum $\sigma_i^{+w_k}$ ($\sigma_i^{-w_k}$). By Lemma 8 the entries of all A_k lie in the interval $[0, \lambda_1 + w]$. But then the sequence $\{A_k\}$ must have a convergent subsequence. Let A_0 be the limit of such a subsequence. Then, by continuity of the spectrum, A_0 has spectrum σ_i^{+w} (σ_i^{-w}). Also, A_0 is a symmetric nonnegative matrix. Therefore, σ_i^{+s} (σ_i^{-s}) is realizable by a

symmetric nonnegative matrix for all $0 \leq s \leq w$. Also, by our assumption, there exists a positive real number s_1 , such that $\sigma_i^{+(w+s)}$ ($\sigma_i^{-(w+s)}$) is realizable by a nonnegative symmetric matrix with Perron eigenvalue $\lambda_1 + w + s$ for all $0 \leq s < s_1$. We conclude that $w + s_1 \in W^+$ ($w + s_1 \in W^-$), but this contradicts the fact that w is the supremum of W^+ (W^-). $\square$

Theorem 8. *Let $\sigma = (\lambda_1, \lambda_2, \dots, \lambda_5)$ be a list of real numbers. Assume σ meets the conditions of Lemma 5. Then σ_i^{-s} is realizable by a 5×5 symmetric nonnegative matrix for any positive real number s and any $i \in \{2, 3, 4, 5\}$.*

Proof. By Theorem 5 we know that σ_i^{-s} is realizable by a 5×5 symmetric nonnegative matrix for sufficiently small positive real number s and any $i \in \{2, 3, 4, 5\}$. We follow the proof of Theorem 7 with the addition of the conditions of Lemma 5, which meets the assumptions of Theorem 7. The delicate step in the proof is when looking at the limit A_0 of a subsequence $\{A_k\}$ of 5×5 symmetric nonnegative matrices, whose spectra meet the conditions of Lemma 5. Let $\tilde{\sigma} = (\delta_1, \delta_2, \dots, \delta_5)$ be the spectrum of A_0 . We want to prove that $\tilde{\sigma}_i^{-s}$ is realizable by a 5×5 symmetric nonnegative matrix for sufficiently small positive real number s and any $i \in \{2, 3, 4, 5\}$. If this can be achieved then the rest of the proof of Theorem 7 is valid as well.

$\tilde{\sigma}$ may not meet some conditions of Lemma 5. The problematic conditions are the strict inequalities $\lambda_5 > -\lambda_1$ and $\lambda_3 > e_1(\sigma)$, which may become an equality in the limit. We look at the different cases:

1. $\delta_5 > -\delta_1$ and $\delta_3 > e_1(\tilde{\sigma})$.

In this case $\tilde{\sigma}_i^{-s}$ is realizable by a 5×5 symmetric nonnegative matrix for sufficiently small positive real number s and any $i \in \{2, 3, 4, 5\}$ by Theorem 5.

2. $\delta_3 = e_1(\tilde{\sigma})$.

In this case $\tilde{\sigma}_i^{-s}$ is realizable by a 5×5 symmetric nonnegative matrix for any positive real number s and any $i \in \{2, 3, 4, 5\}$ by Theorem 4.

3. $\delta_5 = -\delta_1$ and $\delta_3 > e_1(\tilde{\sigma})$.

This case is ruled out by Lemma 6, because we know that $\tilde{\sigma}$ is realizable by the 5×5 symmetric nonnegative matrix A_0 .

$\square$

Theorem 9. *Let $\sigma = (\lambda_1, \lambda_2, \dots, \lambda_5)$ be a list of real numbers and $g \in \mathbb{R}$. Assume σ and g meet the conditions of Theorem 6. Then σ_i^{-s} is realizable by a 5×5 symmetric nonnegative matrix for any positive real number s and any $i \in \{2, 3, 4, 5\}$.*

Proof. By Theorem 6 we know that σ_i^{-s} is realizable by a 5×5 symmetric nonnegative matrix for sufficiently small positive real number s and any $i \in \{2, 3, 4, 5\}$. We follow the proof of Theorem 7 with the addition of the conditions of Theorem 6, which meets the assumptions of Theorem 7. The delicate step in the proof is when looking at the limit A_0 of a subsequence $\{A_k\}$ of 5×5 symmetric nonnegative matrices, whose spectra meet the conditions of Theorem 6 (together with some $g_k \in [0, \frac{1}{2}e_1(\sigma)]$). Let $\tilde{\sigma} = (\delta_1, \delta_2, \dots, \delta_5)$ be

the spectrum of A_0 . We know that the sequence of polynomials $Q_{\sigma_i^{-w_k}}(z)$, each having a real root $g_k \in [0, \frac{1}{2}e_1(\sigma)]$, converges to the polynomial $Q_{\tilde{\sigma}}(z)$. By continuity of roots of polynomials, $Q_{\tilde{\sigma}}(z)$ must have a real root $\tilde{g} \in [0, \frac{1}{2}e_1(\sigma)]$. We want to prove that $\tilde{\sigma}_i^{-s}$ is realizable by a 5×5 symmetric nonnegative matrix for sufficiently small positive real number s and any $i \in \{2, 3, 4, 5\}$. If this can be achieved then the rest of the proof of Theorem 7 is valid as well.

$\tilde{\sigma}$ and $\tilde{g}$ may not meet some conditions of Theorem 6. The problematic conditions are the strict inequalities $\lambda_3 > e_1(\sigma)$ and $r(\sigma) < 0$, which may become an equality in the limit. We look at the different cases:

1. $\delta_3 > e_1(\tilde{\sigma})$ and $r(\tilde{\sigma}) < 0$.

In this case $\tilde{\sigma}_i^{-s}$ is realizable by a 5×5 symmetric nonnegative matrix for sufficiently small positive real number s and any $i \in \{2, 3, 4, 5\}$ by Theorem 6.

2. $\delta_3 = e_1(\tilde{\sigma})$.

In this case $\tilde{\sigma}_i^{-s}$ is realizable by a 5×5 symmetric nonnegative matrix for any positive real number s and any $i \in \{2, 3, 4, 5\}$ by Theorem 4.

3. $\delta_3 > e_1(\tilde{\sigma})$ and $r(\tilde{\sigma}) = 0$.

- (a) $\delta_5 = -\delta_1$.

This case is ruled out by Lemma 6, because we know that $\tilde{\sigma}$ is realizable by the 5×5 symmetric nonnegative matrix A_0 .

- (b) $\delta_5 > -\delta_1$.

In this case $\tilde{\sigma}_i^{-s}$ is realizable by a 5×5 symmetric nonnegative matrix for sufficiently small positive real number s and any $i \in \{2, 3, 4, 5\}$ by Theorem 5.

□

6 Examples

Example 1. Let $\sigma = (1000, 381, 360, -641, -750)$. Then $e_1(\sigma) = 350$ and $r(\sigma) = 306540$, so σ meets the conditions of Lemma 5. Therefore, σ is realizable by the pattern $A(\sigma)$. On the other hand,

$$Q_{\sigma}(z) = 2z^3 + 780z^2 - 169279z - 5139810$$

has approximately the roots $-538.3523722, -27.19321311, 175.5455853$, so it has no root in the range $[0, 175]$. Therefore, σ does not meet the conditions of Lemma 7. Also, as far we know, no other symmetric realizability criterion is satisfied by σ .

Example 2. Let $\sigma = (1000, 370, 367, -637, -750)$. Then $e_1(\sigma) = 350$ and

$$Q_{\sigma}(z) = 2z^3 + 766z^2 - 189010z - 901830$$

has approximately the roots $-552.5556695, -4.683524431, 174.2391939$, so its largest root, g , is in the range $[0, 175]$. Therefore, σ meets the conditions of Lemma 7 and is realizable by the pattern $B(\sigma, g)$. On the other hand, $r(\sigma) = -127980$, so σ does not meet the conditions of Lemma 5. Also, as far we know, no other symmetric realizability criterion is satisfied by σ .

Acknowledgement

I wish to thank Raphi Loewy for his helpful comments after reading an earlier draft of this paper and for suggesting simplified proofs for Lemma 6 and Theorem 4.

References

- [1] A. Berman and R. J. Plemmons. *Nonnegative matrices in the mathematical sciences*, volume 9. Society for Industrial and Applied Mathematics, SIAM; Philadelphia, PA, 1994.
- [2] P. D. Egleston, T. D. Lenker, and S. K. Narayan. The nonnegative inverse eigenvalue problem. *Linear Algebra and its Applications*, 379:475–490, 2004.
- [3] M. Fiedler. Eigenvalues of nonnegative symmetric matrices. *Linear Algebra and its Applications*, 9:119–142, 1974.
- [4] W. Guo. Eigenvalues of nonnegative matrices. *Linear Algebra and its Applications*, 266:261–270, 1997.
- [5] C. R. Johnson, T. J. Laffey, and R. Loewy. The real and the symmetric nonnegative inverse eigenvalue problems are different. *Proceedings of the American Mathematical Society*, 124(12):3647–3651, 1996.
- [6] T. J. Laffey. Perturbing non-real eigenvalues of nonnegative real matrices. *Electronic Journal of Linear Algebra*, 12:73–76, 2004/05.
- [7] R. Loewy and D. London. A note on an inverse problem for nonnegative matrices. *Linear and Multilinear Algebra*, 6(1):83–90, 1978.
- [8] R. Loewy and J. J. McDonald. The symmetric nonnegative inverse eigenvalue problem for 5×5 matrices. *Linear Algebra and its Applications*, 393:275–298, 2004.
- [9] J. J. McDonald and M. Neumann. The soules approach to the inverse eigenvalue problem for nonnegative symmetric matrices of order $n \leq 5$. *Contemporary Mathematics*, 259:387–407, 2000.
- [10] O. Rojo and R. L. Soto. Guo perturbation for symmetric nonnegative circulant matrices. *Linear Algebra and its Applications*, 431(5-7):594–607, 2009.
- [11] O. Spector. A characterization of trace zero symmetric nonnegative 5×5 matrices. *Linear Algebra and its Applications*, 434(4):1000–1017, 2011.
- [12] H. R. Suleimanova. Stochastic matrices with real characteristic values. *Doklady Akademii nauk SSSR*, 66:343–345, 1949.